



ISSN (P): 2521-3466
ISSN (E): 2521-3474
© Clinical Orthopaedics
www.orthoresearchjournal.com
2024; 8(3): 35-36
Received: 16-07-2024
Accepted: 23-08-2024

Dr. Siddaram Patil N
Professor & H.O.D, Department
of Orthopaedics, Dr. PMR
Institute of medical Sciences &
Teaching Hospital, Chevella,
RR District, Telangana, India

Dr. Azam
Assistant Professor, Department
of Orthopaedics, Dr. PMR
Institute of medical Sciences &
Teaching Hospital, Chevella,
RR District, Telangana, India

Dr. Sai Tej V
Junior Resident, Department of
Orthopaedics, Dr. PMR Institute
of medical Sciences & Teaching
Hospital, Chevella, RR District,
Telangana, India

Corresponding Author:
Dr. Azam
Assistant Professor, Department
of Orthopaedics, Dr. PMR
Institute of medical Sciences &
Teaching Hospital, Chevella,
RR District, Telangana, India

A comparative prospective study of platelet rich plasma vs corticosteroid injection in lateral epicondylitis- research article

Dr. Siddaram Patil N, Dr. Azam and Dr. Sai Tej V

DOI: <https://doi.org/10.33545/orthor.2024.v8.i3a.451>

Abstract

Introduction: Lateral epicondylitis, commonly referred to as 'tennis elbow' is seen to affect 1% to 3% of the general population in the 3rd and 4th decade of life. Treatment modalities for the treatment of lateral epicondylitis include analgesics and immobilisation. 90% of cases resolve spontaneously within 6-12 months.

Materials and Methods: Prospective comparative study performed at department of orthopaedics in Dr PMR institute of Medical Sciences from September 2023 to November 2024, 20 patients (12 females and 8 males) were included in the study after explaining the procedure with consent.

Results: We included 20 patients in this study, 12 were females and 8 were males with chronic pain at elbow for more than 3 months not responding to 6 weeks of conservative management. Patient follow up was done at fourth, eighth and twelfth week of post injection. Mean VAS & DASH pre injection score average for platelet rich plasma group was 66.6 & 56.4 respectively while 36.5 & 29.1 at the end of one year treatment. Mean VAS & DASH pre injection score average for steroid group was 65.5 & 55.2 respectively while 37.8 & 34.0 at the end of one year treatment.

Conclusion: A single injection of autologous platelet rich plasma improves elbow pain and functional activities more effectively than corticosteroid injection in lateral epicondylitis.

Keywords: Tennis elbow, platelet rich plasma, corticosteroid, lateral epicondylitis

Introduction

Lateral epicondylitis, commonly referred to as 'tennis elbow' is seen to affect 1% to 3% of the general population in the 3rd and 4th decade of life. It is one of the most common causes for elbow pain. The pathogenesis of an overuse injury is thought to be a result from cumulative micro trauma that weakens the structural and vascular elements of the tendon. Micro trauma to a mechanical structure occurs even if the loads are within the material's strength limits, and is due to fatigue after repetitive loads. If the muscle is weak or fatigued, the energy absorbing capacity of the whole muscle-tendon unit is reduced, and tendon stresses will increase. Treatment modalities for the treatment of lateral epicondylitis include analgesics and immobilisation. 90% of cases resolve spontaneously within 6-12 months. Other modalities include wrist bracing, elbow bracing, local corticosteroid injection, shockwave therapy and modifying poor technique in sport or work have been used in the treatment for Lateral Epicondylitis.

Materials and Methods

Prospective comparative study performed at department of orthopaedics in Dr PMR Institute of Medical Sciences from September 2023 to November 2024, 20 patients (12 females and 8 males) were included in the study after explaining the procedure with consent by block randomization with 20 in each group.

Inclusion criteria

Patients in the age group of 25-50 years

Patients with symptoms of lateral epicondylitis for more than 3 months not responding to 6 weeks of conservative management.

Exclusion criteria

Diabetic patients: Patients suffering with rheumatoid arthritis. Patients with history of trauma or surgery to the lateral epicondyle. Patients who have previously received an injection at the lateral epicondyle in the last 3 months. Patients suspicion of nerve involvement.

Methodology

1ml (40 mg) of Triamcinolone acetonide was taken with 2ml of Lignocaine (1%). It was injected in the maximum tenderness point deep into the tendon. The patient was then observed for 15 to 20 minutes and then discharged. The platelet rich plasma preparation has been done using desktop from the patient with 18 gauge needle.

Blood is equally divided into three parts (nine ml each) which is then added to three pre filled test tubes, each containing one ml of 3.8% of sodium citrate solution. The blood is centrifuged at 1500 rpm for 15 minutes. By the end of the procedure the whole blood is separated into three layers such as platelet poor plasma (PPP), platelet rich plasma (PRP) and red blood cells (RBC). Platelet rich plasma is withdrawn from the middle layer and 3-4 ml of platelet rich plasma was prepared

PRP Injection Technique

The procedure was done on an outpatient basis. Once the exact location was determined by assessing the maximum tenderness point clinically, the patient was injected with a local anesthetic drug (Lignocaine) under sterile technique.

Platelet rich plasma was injected with 3-4 ml platelet rich plasma, using a “peppering” technique in a clock wise manner to better cover the affected area of lateral epicondyle.

Post injection protocol

Since the patients may experience discomfort at the site of the injection for up to three days, they are advised to have ice fermentation over the injection site, limb elevation, activity modification and oral acetaminophen for pain relief.

Follow UP

All the patients were followed up at fourth, eighth and twelfth week of post injection.

One patient did not return for final follow up in platelet rich plasma (PRP) group. At follow up, pain was assessed according to Visual Analogue Scale (VAS) and Disabilities of the Arm, Shoulder and Hand (DASH) score and compared with pre injection score levels. Final outcome was measured based on the pain reduction from the pre injection level. Patients were observed for post.

Results

The average Visual Analogue Scale (VAS) and Disabilities of the Arm, Shoulder and Hand (DASH) scores in both the groups of pre injection, four, eight and 12 weeks post injection are shown in the tables:

For Platelet Rich Plasma Group

- The visual analogue scale (VAS) score is decreased by 30.1
- And the disability of arm, shoulder and hand (DASH) score is decreased by 27.3 at 12 weeks compared to the pre injection score.

For Corticosteroid Group

- The visual analogue scale (VAS) score is decreased by 27.7
- And the disabilities of the arm, shoulder and hand (DASH)

score is decreased by 21.2 at 12 weeks compared to the pre injection score.

Complications

In steroid group one patient had paraesthesia at the injection site of elbow at fourth week post injection but it disappeared at twelfth week with observation. No case of infection, cellulitis was observed. No neurovascular injury noted.

Discussion

Nirschl observed that the basic pathology was in the origin of the extensor carpi radialis brevis (ECRB) tendon. But sometimes the anteromedial edge of extensor digitorum communis (EDC) and the deep surface of extensor carpiradialis longus (ECRL) may also be involved. In the study by Gosen *et al.* march 2011, compared the effectiveness of autologous platelet rich plasma injection to steroid injection therapy in lateral epicondylitis, it is proved that platelet rich plasma injection is safe and easy. Concerning functional impairment, the corticosteroid group showed better results during the initial period and then declined to baseline level. Whereas in platelet rich plasma group symptoms improved progressively. In the Mishra and Pavelko research, the success rate was 93% in the platelet rich plasma group and 65% success rate for steroid group in the Hay *et al.* study.

Conflict of Interest: Not available

Financial Support: Not available

References

1. Thanasas C, Papadimitriou G, Charalambidis C, Paraskevopoulos I, Papanikolaou A. Platelet-rich plasma Vs autologous whole blood for treatment of chronic lateral elbow epicondylitis: ARCT. *Am J Sports Med.* 2011;39:2130-2134.
2. Shiri R, Viikari-juntura E, Vareven H, Heliövaara M. Prevalence and determinants of lateral and medial epicondylitis: A population study. *Am J Epidemiology.* 2006;164(11):1065-1074.
3. Zeisig E, Fahlstrom M, Ohberg L, Alfredson H. A 2 – year sonographic follow-up after intratendinous injection therapy in patients with lateral epicondylitis. *Br J Sports Med.* 2008;42(4):267-71.
4. Peerbooms JC, Sluimer J, Brujin DJ, Gosens T. Positive effect of an autologous platelet concentrate in lateral epicondylitis in a double-blind randomized controlled trial: platelet-rich plasma versus corticosteroid injection with a 1-year follow-up. *Am J Sports Med.* 2010;38(2):255-262.
5. Sampson S, Gerhardt M, Mandelbaum B. Platelet rich plasma injection grafts for musculoskeletal injury. *Cur rev of musculoskeletal med.* 2008;1:165-174.

How to Cite This Article

Patil SN, Azam, Tej SV. A comparative prospective study of platelet rich plasma vs corticosteroid injection in lateral epicondylitis-research article. *National Journal of Clinical Orthopaedics.* 2024;8(3):35-36.

Creative Commons (CC) License

This is an open-access journal, and articles are distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 International (CC BY-NC-SA 4.0) License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.